

**Chapter 6, Table T9.** Occurrence of selected benthic foraminifer taxa, Site 1265.

Notes: Occurrence: C = common, F = few, R = rare. Preservation: E = excellent, G = good, M = moderate, P = poor. Reworking: R = reworking and downslope transport, R? = reworking and downslope transport probable. Paleodepth: UA = upper abyssal, LB = lower bathyal, DT = downslope transport, ? = unknown. x = present, xx = dominant species, \* = reworked.

**Chapter 6, Table T9.** Occurrence of selected benthic foraminifer taxa, Site 1265. (See table notes. Continued on next five pages.)

Table T9 (continued).

**Table T9 (continued).**

**Table T9 (continued).**

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Abundance	Preservation	Reworking	Paleodepth	<i>Abyssamina poagi</i>	<i>Abyssamina quadrata</i>	<i>Alabamina creta</i>	<i>Alabamina dissonata</i>	<i>Alabaminella weddellensis</i>	<i>Anomaliodes rubiginosa</i>	<i>Anomaliodes semicirratus</i>	<i>Anomaliodes spissiformis</i>	<i>Aragonia aragonensis</i>	<i>Aragonia velascoensis</i>	<i>Bigennerina nodosaria</i>	<i>Bolivina</i> sp. (small)	<i>Bolivinita pseudothalmani</i>	<i>Bolivinoides delicatulus</i>	<i>Bolivinoides huneri</i>	<i>Buliminella elongata</i>	<i>Buliminella exilis</i>	<i>Buliminella kugleri</i>	<i>Buliminella rostrata</i>	<i>Buliminella semicostata</i>	<i>Buliminella simplex</i>	<i>Buliminella thanensis</i>	<i>Buliminella trinitatis</i>	<i>Buliminella turopensis</i>	<i>Buliminella velascoensis</i>	<i>Buliminella</i> sp.	<i>Cassidulina laevigata</i>	<i>Cibicidoides grimsdalei</i>	<i>Cibicidoides hyphalus</i>
1265D-2X-CC, 23–28	267.33	305.60	R	G	?	x																													
1265A-28H-CC, 17–22	266.46	305.91	F	G	?	x																													
1265D-3X-CC, 19–24	272.57	311.26	F	G	?	x	x																												
1265A-29H-6, 148–149	274.36	315.14	F	G	?	x	x																												
1265A-29H-7, 8–9	274.46	315.24	F	M	?	x	x																												
1265A-29H-7, 30–31	274.68	315.46	R	M	?	x	x																												
1265D-4H-CC, 5–7	274.82	315.65	F	M	?	x	x																												
1265A-29H-7, 50–51	274.88	315.66	F	M	?	x	x																												
1265D-4H-CC, 27–28	274.88	315.71	F	M	?	x	x																												
1265A-29H-7, 65–66	275.03	315.81	F	M	?	xx																													
1265A-29H-7, 70–71	275.08	315.86	R	G	LB	x	x																												
1265D-5H-CC, 9–10	274.89	315.87	R	G	LB	x	x																												
1265A-29H-7, 80–81	275.18	315.96	R	G	LB	x	x																												
1265A-29H-7, 128–129	275.66	316.44	R	G	LB	x	x																												
1265A-29H-CC, 28–33	276.16	316.94	R	G	LB	x	x																												
1265A-30H-CC, 15–20	276.68	318.79	R	G	LB	x	x																												
1265A-31H-CC, 32–37	278.21	320.49	R	G	LB	x	x																												
1265A-32H-CC, 16–21	285.56	328.05	R	G	LB	x	x																												
1265A-34X-CC, 38–43	296.21	340.64	R	G	LB	x	x																												
1265A-35X-CC, 30–35	309.20	355.04	R	G	LB	x	x																												
1265A-36X-CC, 35–40	312.36	359.48	G	LB	x	x																													

Notes: Occurrence: C = common, F = few, R = rare. Preservation: E = excellent, G = good, M = moderate, P = poor. Reworking: R = reworking and downslope transport, R? = reworking and

**Table T9 (continued).**

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Abundance	Preservation	Reworking	Paleodepth	<i>Cibicidoides mundulus</i>	<i>Cibicidoides eoceneus</i>	<i>Cibicidoides praemundulus</i>	<i>Cibicidoides velascoensis</i>	<i>Cibicidoides wuellestorfi</i>	<i>Clavulinoides</i> spp.	<i>Clinapertina complanata</i>	<i>Clinapertina inflata</i>	<i>Clinapertina subplanispira</i>	<i>Conophistoma midwayensis</i>	<i>Eggerella bradyi</i>	<i>Epistomina exigua</i>	<i>Furstenkoia</i> sp.	<i>Gaudryina laevigata</i>	<i>Gaudryina pyramidata</i>	<i>Globocassidulina subglobosa</i>	<i>Gyroidinoides beisseli</i>	<i>Gyroidinoides globosus</i>	<i>Gyroidinoides</i> spp.	<i>Hoeglundina elegans</i>	<i>Karreniella bradyi</i>	<i>Karreniella subglabra</i>	<i>Laevidentalina</i> spp.	<i>Laticarinina pauperata</i>	<i>Lenticulina</i> spp.	<i>Marssonella oxycona</i>	<i>Melonis</i> spp.	Miliolids	<i>Norion havanense</i>	<i>Nuttallides umbonifera</i>
1265D-2X-CC, 23–28	267.33	305.60	R G	?	x																															
1265A-28H-CC, 17–22	266.46	305.91	F G	?	x																															
1265D-3X-CC, 19–24	272.57	311.26	F G	?																																
1265A-29H-6, 148–149	274.36	315.14	F G	?																																
1265A-29H-7, 8–9	274.46	315.24	F G	?																																
1265A-29H-7, 30–31	274.68	315.46	R M	?																																
1265D-4H-CC, 5–7	274.82	315.65	F M	?																																
1265A-29H-7, 50–51	274.88	315.66	F M	?																																
1265D-4H-CC, 27–28	274.88	315.71	F M	?																																
1265A-29H-7, 65–66	275.03	315.81	F M	?																																
1265A-29H-7, 70–71	275.08	315.86	R G	LB			x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x						
1265D-5H-CC, 9–10	274.89	315.87	R G	LB			x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x						
1265A-29H-7, 80–81	275.18	315.96	R G	LB			x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-29H-7, 128–129	275.66	316.44	R G	LB			x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-29H-CC, 28–33	276.16	316.94	R G	LB			x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-30H-CC, 15–20	276.68	318.79	R G	LB	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-31H-CC, 32–37	278.21	320.49	R G	LB			x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-32H-CC, 16–21	285.56	328.05	R G	LB			x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-34X-CC, 38–43	296.21	340.64	R G	LB			x		x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-35X-CC, 30–35	309.20	355.04	R G	LB	x	x			x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							
1265A-36X-CC, 35–40	312.36	359.48	G	LB	x	x			x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x							

downslope transport probable. Paleodepth: UA = upper abyssal, LB = lower bathyal, DT = downslope transport, ? = unknown. x = present, xx = dominant species, \* = reworked.

**Table T9 (continued).**

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Abundance	Preservation	Reworking	Paleodepth	<i>Nuttallides truempyi</i>	<i>Nuttallella florealis</i>	<i>Oridorsalis umbonatus</i>	<i>Orthomorphina</i> spp.	<i>Oscularia velscoensis</i>	<i>Paralabamina</i> spp.	<i>Polymorphinid</i> taxa	<i>Plectofrondiculata paucicostata</i>	<i>Pleurostomellid</i> taxa	<i>Pullenia conyelli</i>	<i>Pullenia jarvisi</i>	<i>Pullenia</i> spp.	<i>Quadrimorphina</i> spp.	<i>Rectabulimina carpenterae</i>	<i>Sigmilospsis schumbergeri</i>	<i>Siphogenerinoides brevispinosa</i>	<i>Siphonodosaria hispidula</i>	<i>Siphonodosaria lepidula</i>	<i>Siphonodosaria</i> spp.	<i>Spiroplectammina spectabilis</i>	<i>Stainforthia complanata</i>	<i>Stensioeina beccaniformis</i>	<i>Tappanina salmensis</i>	<i>Tritaxia havanensis</i>	Unilocular taxa	<i>Uvigerina graciiformis</i>	<i>Uvigerina peregrina</i> group	<i>Vulvulina spinosa</i>
1265D-2X-CC, 23–28	267.33	305.60	R	G	?	x	x	x	x									x																
1265A-28H-CC, 17–22	266.46	305.91	F	G	?	x	x	x	x									x																
1265D-3X-CC, 19–24	272.57	311.26	F	G	?	x	x	x										x																
1265A-29H-6, 148–149	274.36	315.14	F	G	?	xx	x	x	x								x*	x	x															
1265A-29H-7, 8–9	274.46	315.24	F	G	?	xx	x	x	x								x	x	xx															
1265A-29H-7, 30–31	274.48	315.46	R	M	?	xx	x	x	x					x			x	x	x															
1265D-4H-CC, 5–7	274.82	315.65	F	M	?	xx	x	x	x				x			x	x	x																
1265A-29H-7, 50–51	274.88	315.66	F	M	?	xx	x	x	x				x			x	x	x																
1265D-4H-CC, 27–28	274.88	315.71	F	M	?	xx	xx	x	x				x			x	x	x																
1265A-29H-7, 65–66	275.03	315.81	F	M	?	x	x	x	x				x			x	x	x																
1265A-29H-7, 70–71	275.08	315.86	R	G	LB	x	x	x	x				x	x		x	x	x	x	x														
1265D-5H-CC, 9–10	274.89	315.87	R	G	LB	x	x	x	x				x	x		x	x	x	x	x														
1265A-29H-7, 80–81	275.18	315.96	R	G	LB	x	x	x	x				x	x		x	x	x	x	x														
1265A-29H-7, 128–129	275.66	316.44	R	G	LB	x	x	x	x				x	x		x	x	x	x	x														
1265A-29H-CC, 28–33	276.16	316.94	R	G	LB	x	x	x	x				x	x		x	x	x	x	x											x			
1265A-30H-CC, 15–20	276.68	318.79	R	G	LB	x	x	x	x				x	x		x	x	x	x	x										x	x			
1265A-31H-CC, 32–37	278.21	320.49	R	G	LB	x	x	x	x	x			x	x		x	x	x	x	x														
1265A-32H-CC, 16–21	285.56	328.05	R	G	LB	x	x	x	x				x	x		x	x	x	x	x														
1265A-34X-CC, 38–43	296.21	340.64	R	G	LB	x	x	x	x				x	x		x	x	x	x	x														
1265A-35X-CC, 30–35	309.20	355.04	R	G	LB	x	x	x	x	x			x	x		x	x	x	x	x										x				
1265A-36X-CC, 35–40	312.36	359.48		G	LB	x	x	x	x	x			x	x		x	x	x	x	x										x				