





1084A-1H 0.0-3.0 mbsf										
Leg 175 Site 1084 Hole A Core 1H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
0.0										DIATOM-RICH CLAYEY NANNOFOSSIL OOZE and FORAMINIFER-BEARING DIATOM-RICH CLAYEY NANNOFOSSIL OOZE. The core consists of olive (5Y 5/3) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE and dark olive gray (5Y 3/2) FORAMINIFER-BEARING DIATOM-RICH CLAYEY NANNOFOSSIL OOZE.
1.0									SS	
2.0									SS	
3.0									IW	
									PAL	




1084A-2H 3.0-12.5 mbsf										
Leg 175 Site 1084 Hole A Core 2H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY
2									SS	The core consists of homogeneous dark olive gray (5Y 3/2) FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY.
3									IW	
4										
5									SS	
6									IW	
7									SS	
8										
9									PAL	



1084A-2H


1084A-3H 12.5-22.0 mbsf										
Leg 175 Site 1084 Hole A Core 3H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1	1									FORAMINIFER-BEARING NANNOFOSSIL CLAY, FORAMINIFER- AND DIATOM-BEARING NANNOFOSSIL CLAY and NANNOFOSSIL- AND DIATOM-BEARING CLAY
2	2								SS	The core consists of olive (5Y 4/3) FORAMINIFER-BEARING NANNOFOSSIL CLAY from Section 1 to Section 3, olive (5Y 5/4) FORAMINIFER- AND DIATOM-BEARING NANNOFOSSIL CLAY from Section 4 to Section 5, 60 cm, and from Section 7 to the end of the core. Section 5, 60 cm, to Section 6 is dark olive brown (5Y 3/3) NANNOFOSSIL- AND DIATOM-BEARING CLAY.
3	3									
4	4								IW	
5	5								SS	
6	6									
7	7									
8	8								SS	
10	10								PAL	





1084A-4H 22.0-31.5 mbsf										
Leg 175 Site 1084 Hole A Core 4H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-BEARING NANNOFOSSIL-RICH CLAY, DIATOM-BEARING FORAMINIFER-RICH CLAY, and NANNOFOSSIL- AND FORAMINIFER-BEARING DIATOM-RICH CLAY The core consists of dark olive gray (5Y 3/2) DIATOM-BEARING NANNOFOSSIL-RICH CLAY, very dark grayish brown NANNOFOSSIL- AND FORAMINIFER-BEARING DIATOM-RICH CLAY, and olive NANNOFOSSIL- AND FORAMINIFER-BEARING DIATOM-RICH CLAY. Section 1, 0-100 cm, is mottled, and contains several very dark grayish brown clasts. Section 6 contains abundant diatom resting spores.
2									IW	
3									SS	
4										
5										
6										
7									SS	
8										
9										
10									PAL	















1084A-8H 60.0-69.5 mbsf								
Leg 175 Site 1084 Hole A Core 8H								
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DESCRIPTION
1								FORAMINIFER-BEARING CLAY-RICH NANNOFOSSIL OOZE and DIATOM-BEARING CLAY-RICH NANNOFOSSIL OOZE The core consists of olive (5Y 5/3) DIATOM-BEARING CLAY-RICH NANNOFOSSIL OOZE. Section 1, 110 cm, to Section 3, 10 cm, is olive gray (5Y 5/2) FORAMINIFER-BEARING CLAY-RICH NANNOFOSSIL OOZE.
2								
3								
4								
5								
6								
7								
8								
9								
10								

1084A-8H













1084A-13H 107.5-117.0 mbsf									
Leg 175 Site 1084 Hole A Core 13H									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									
2									SS
3									
4									
5									SS
6									
7									
8									PAL
10									


~ NANNOFOSSIL-RICH DIATOMACEOUS CLAY and NANNOFOSSIL DIATOMACEOUS CLAY

The core consists of olive (5Y 4/3) NANNOFOSSIL-RICH DIATOMACEOUS CLAY which grades into NANNOFOSSIL DIATOMACEOUS CLAY.








1084A-16H 136.0-145.5 mbsf										
Leg 175 Site 1084 Hole A Core 16H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										NANNOFOSSIL-AND DIATOM-RICH CLAY and DIATOM-RICH NANNOFOSSIL CLAY The core consists of dark olive gray (5Y 3/2) NANNOFOSSIL-AND DIATOM-RICH CLAY. Section 4, 30-150 cm, is olive (5Y 5/3) DIATOM-RICH NANNOFOSSIL CLAY.
2								SS		
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										
26										
27										
28										
29										
30										
31										
32										
33										
34										
35										
36										
37										
38										
39										
40										
41										
42										
43										
44										
45										
46										
47										
48										
49										
50										
51										
52										
53										
54										
55										
56										
57										
58										
59										
60										
61										
62										
63										
64										
65										
66										
67										
68										
69										
70										
71										
72										
73										
74										
75										
76										
77										
78										
79										
80										
81										
82										
83										
84										
85										
86										
87										
88										
89										
90										
91										
92										
93										
94										
95										
96										
97										
98										
99										
100										
101										
102										
103										
104										
105										
106										
107										
108										
109										
110										
111										
112										
113										
114										
115										
116										
117										
118										
119										
120										
121										
122										
123										
124										
125										
126										
127										
128										
129										
130										
131										
132										
133										
134										
135										
136										
137										
138										
139										
140										
141										
142										
143										
144										
145										
146										
147										
148										
149										
150										
151										
152										
153										
154										
155										
156										
157										
158										
159										
160										
161										
162										
163										
164										
165										
166										
167										
168										
169										
170										
171										
172										
173										
174										
175										
176										
177										
178										
179										
180										
181										
182										
183										
184										
185										
186										
187										
188										
189										
190										
191										
192										
193										
194										
195										
196										
197										
198										
199										
200										
201										
202										
203										
204										
205										
206										
207										
208										
209										
210										
211										
212										
213										
214										
215										
216										
217										
218										
219										
220										
221										
222										
223										
224										
225										
226										
227										
228										
229										
230										
231										
232										
233										
234										
235									</	

1084A-16H


1084A-17H 145.5-149.5 mbsf										
Leg 175 Site 1084 Hole A Core 17H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-RICH NANNOFOSSIL CLAY, and CLAYEY DIATOMACEOUS NANNOFOSSIL OOZE The core consists of DIATOM-RICH NANNOFOSSIL CLAY, and CLAYEY DIATOMACEOUS NANNOFOSSIL OOZE. Section 1, 0-100 cm, is dark olive gray (5Y 3/2) DIATOM-RICH NANNOFOSSIL CLAY. Section 1, 100 cm, to the end of the core is pale olive (5Y 6/3) CLAYEY DIATOMACEOUS NANNOFOSSIL OOZE.
2									SS	
3										
4									PAL	



1084A-18X 149.5-152.5 mbsf									
Leg 175 Site 1084 Hole A Core 18X									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
DOLOMITE HORIZON									
The core consists of light gray (5Y 6/1) DOLOMITE HORIZON.									




1084A-19X 152.5-162.2 mbsf										
Leg 175 Site 1084 Hole A Core 19X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								///		CLAYEY DIATOMACEOUS NANNOFOSSIL OOZE The core consists of homogeneous olive (5Y 4/3) CLAYEY DIATOMACEOUS NANNOFOSSIL OOZE.
2								///	SS	
3								///	IW	
4								///	SS	
6								///	PAL	



1084A-20X 162.2-171.8 mbsf										
Leg 175 Site 1084 Hole A Core 20X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5 6 7									— SS — SS — SS — SS — PAL	NANNOFOSSIL-BEARING DIATOM-RICH CLAY, NANNOFOSSIL-RICH DIATOMACEOUS CLAY, and DIATOMACEOUS CLAYEY NANNOFOSSIL OOZE The core consists of dark olive gray (5Y 3/2) and olive brown (2.5Y 3/3) NANNOFOSSIL-BEARING DIATOM-RICH CLAY from Section 1 to Section 3, 40 cm, olive (5Y 4/3) NANNOFOSSIL-RICH DIATOMACEOUS CLAY from Section 3, 40 cm to Section 4, 110 cm, olive brown (2.5Y 3/3) NANNOFOSSIL-BEARING DIATOM-RICH CLAY from Section 4, 110 cm, to Section 5, 100 cm, and olive gray (5Y 5/2) DIATOMACEOUS CLAYEY NANNOFOSSIL OOZE to the end of the core.

1084A-20X



1084A-21X 171.8-181.5 mbsf										
Leg 175 Site 1084 Hole A Core 21X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY, DIATOMACEOUS NANNOFOSSIL CLAY, and DIATOM-RICH CLAYEY NANNOFOSSIL OOZE The core consists of olive gray (5Y 4/2) FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY from Section 1 to Section 2, olive gray (5Y 4/2) DIATOMACEOUS NANNOFOSSIL CLAY from Section 3 to Section 4, olive (5Y 4/3) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE in Section 5, and black (5Y 2.5/1) FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY in Section 6 to the end of the core.
2									SS	
3									SS	
4									PAL	
5									SS	
6									SS	
7									PAL	




1084A-22X 181.5-191.2 mbsf									
Leg 175 Site 1084 Hole A Core 22X									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
181.5									SS
191.2									PAL
The core consists of black (2.5Y 2.5/1) DIATOM-RICH CLAY.									

1084A-22X





1084A-23X 191.2-200.8 mbsf										
Leg 175 Site 1084 Hole A Core 23X										
MEETRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4										CLAYEY NANNOFOSSIL OOZE The core consists of olive (5Y 4/3) CLAYEY NANNOFOSSIL OOZE.



1084A-24X 200.8-210.4 mbsf										
Leg 175 Site 1084 Hole A Core 24X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-RICH CLAYEY NANNOFOSSIL OOZE, DIATOMACEOUS CLAYEY NANNOFOSSIL OOZE, and NANNOFOSSIL- AND DIATOM-RICH CLAY The core consists of olive gray (5Y 4/2) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE from Section 1 to Section 2, olive (5Y 4/3) DIATOMACEOUS CLAYEY NANNOFOSSIL OOZE from Section 3 to Section 4, 70 cm, and black (5Y 2.5/2) NANNOFOSSIL- AND DIATOM-RICH CLAY from Section 4, 70 cm, to the end of the core.
2									SS	
3									SS	
4									SS	
5									SS	
6									PAL	






1084A-26X 220.1-229.7 mbsf										
Leg 175 Site 1084 Hole A Core 26X										
MEETRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
220.1 220.2 220.3 220.4	1 2 3								— SS — IW — SS — PAL	DIATOM-BEARING NANNOFOSSIL-RICH CLAY The core consists of dark olive gray (5Y 3/2) DIATOM-BEARING NANNOFOSSIL-RICH CLAY.




1084A-27X 229.7-239.3 mbsf										
Leg 175 Site 1084 Hole A Core 27X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING FORAMINIFER-NANNOFOSSIL OOZE, DIATOM- AND NANNOFOSSIL-RICH CLAY, and DIATOM-RICH NANNOFOSSIL OOZE The core consists of dark olive gray (5Y 3/2) DIATOM-BEARING FORAMINIFER-NANNOFOSSIL OOZE and DIATOM- AND NANNOFOSSIL-RICH CLAY. Section 4, 60 cm, to Section 5, 150 cm, is olive (5Y 4/3) DIATOM-RICH NANNOFOSSIL OOZE.
2									SS	
3									SS	
4									SS	
5									SS	
6									PAL	



1084A-28X 239.3-248.9 mbsf										
Leg 175 Site 1084 Hole A Core 28X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM- AND NANNOFOSSIL-RICH CLAY and DIATOM-RICH NANNOFOSSIL OOZE The core consists of olive (5Y 4/3) to dark olive gray (5Y 3/2) DIATOM-RICH NANNOFOSSIL OOZE. Section 1, 0-11 cm, is black (5Y 2.5/2) DIATOM- AND NANNOFOSSIL-RICH CLAY. A gradational color change from olive to dark olive gray is present over 40 cm between Section 2 and Section 3.
2									SS	
3										
4										
5										
6									SS	
7										
8										
9										
10									PAL	








1084A-29X 248.9-258.5 mbsf										
Leg 175 Site 1084 Hole A Core 29X										
MEETRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5									SS IW SS PAL	DIATOM-BEARING CLAY and NANNOFOSSIL-BEARING DIATOM-RICH CLAY The core consists of black (5Y 2.5/1) DIATOM-BEARING CLAY which grades over an interval of about 60 cm to very dark gray (5Y 3/1) NANNOFOSSIL-BEARING DIATOM-RICH CLAY in Sections 3 and 4.



1084A-30X 258.5-268.2 mbsf										
Leg 175 Site 1084 Hole A Core 30X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	NANNOFOSSIL- AND DIATOM-BEARING CLAY and DIATOM-BEARING NANNOFOSSIL-RICH CLAY The core consists of very dark gray (5Y 3/1) to olive (5Y 4/3) or very dark grayish brown (2.5Y 3/2) NANNOFOSSIL- AND DIATOM-BEARING CLAY and DIATOM-BEARING NANNOFOSSIL-RICH CLAY.
2									SS	
3									SS	
4								~		
5								~		
6								~	PAL	

1084A-30X

1084A-31X 268.2-277.8 mbsf										
Leg 175 Site 1084 Hole A Core 31X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1	1									FOARAMINIFER- AND DIATOM-BEARING NANNOFOSSIL CLAY, and CLAY-RICH DIATOMACEOUS NANNOFOSSIL OOZE The core consists of olive (5Y 4/3) FORAMINIFER- AND DIATOM-BEARING NANNOFOSSIL CLAY. Section 1, 70 cm, to Section 1, 140 cm, is dark olive gray (5Y 3/2) CLAY-RICH DIATOMACEOUS NANNOFOSSIL OOZE.
2	2								SS	
3	3								SS	
4	4								PAL	




1084A-32X 277.8-287.4 mbsf										
Leg 175 Site 1084 Hole A Core 32X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-RICH NANNOFOSSIL CLAY The core consists of olive gray (5Y 4/2) DIATOM-RICH NANNOFOSSIL CLAY.
2										
3										
4									IW	
4									SS	
5									PAL	

1084A-32X





1084A-33X 287.4-297.0 mbsf										
Leg 175 Site 1084 Hole A Core 33X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								~		DIATOM-RICH NANNOFOSSIL CLAY The core consists of olive gray (5Y 4/2) DIATOM-RICH NANNOFOSSIL CLAY.
2									SS	
3										
4										
5									SS	
6								~	PAL	




1084A-34X 297.0-306.6 mbsf										
Leg 175 Site 1084 Hole A Core 34X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-BEARING NANNOFOSSIL-RICH CLAY, FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY and DIATOM-BEARING NANNOFOSSIL CLAY The core consists of intercalated intervals of olive (5Y 4/3) DIATOM-BEARING NANNOFOSSIL-RICH CLAY and dark olive gray (5Y 3/2) FORAMINIFER-BEARING DIATOM-RICH NANNOFOSSIL CLAY and DIATOM-BEARING NANNOFOSSIL CLAY. Dark olive intervals extend from Section 1, 0-100 cm, Section 3, 110 cm to Section 5, 10 cm, Section 5, 96-130 cm, and the core catcher. Olive intervals extend from Section 1, 100 cm, to Section 3, 110 cm, Section 5, 10-96 cm, and Section 5, 130-146 cm.
2										
3									SS	
4										
5									SS	
6									PAL	




1084A-35X 306.6-316.3 mbsf										
Leg 175 Site 1084 Hole A Core 35X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING NANNOFOSSIL CLAY and DIATOMACEOUS NANNOFOSSIL CLAY The core consists of dark olive gray (5Y 3/2) DIATOMACEOUS NANNOFOSSIL CLAY. Section 2, 10- 80 cm, is dark olive gray (5Y 3/2) DIATOM-BEARING NANNOFOSSIL CLAY.
2								SS		
3								IW		
4								SS		
5								PAL		



1084A-36X 316.3-325.9 mbsf										
Leg 175 Site 1084 Hole A Core 36X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING NANNOFOSSIL CLAY and FORAMINIFER-BEARING DIATOMACEOUS CLAYEY NANNOFOSSIL OOZE The core consists of intercalated intervals of olive (5Y 4/3) DIATOM-BEARING NANNOFOSSIL CLAY and dark olive gray (5Y 3/2) FORAMINIFER-BEARING DIATOMACEOUS CLAYEY NANNOFOSSIL OOZE. The olive colored intervals occur at Section 1, 90 cm, to Section 3, 65 cm, and Section 4, 15 cm, to the end of the core.
2									SS	
3									SS	
4									PAL	





1084A-37X 325.9-335.6 mbsf										
Leg 175 Site 1084 Hole A Core 37X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOMACEOUS CLAY The core consists of dark olive gray (5Y 3/2) and olive (5Y 4/3) DIATOMACEOUS CLAY. Alternating 1-3 cm thick dark olive gray and olive colored bands are present in Section 4, 45-100 cm, Section 5, 5-130 cm, Section 6, 15-45 cm, 70-80 cm, and Section 7, 8-25 cm.
2									SS	
3										
4										
5									SS	
6										
7										
8									PAL	




1084A-38X 335.6-345.2 mbsf										
Leg 175 Site 1084 Hole A Core 38X										
MEETERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								↕	SS	DIATOMACEOUS CLAY and NANNOFOSSIL DIATOMACEOUS CLAY The core consists of DIATOMACEOUS CLAY and NANNOFOSSIL DIATOMACEOUS CLAY. Dark olive gray (5Y 3/2) DIATOMACEOUS CLAY is present from top of the core to Section 2, 100 cm. The other sections are olive (5Y 4/3 and 5Y 4/4) NANNOFOSSIL DIATOMACEOUS CLAY.
2										
3									SS	
4									IW	
5										
6								↕	SS	
7									PAL	




1084A-39X 345.2-354.8 mbsf										
Leg 175 Site 1084 Hole A Core 39X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOMACEOUS CLAY and NANNOFOSSIL-RICH DIATOMACEOUS CLAY The core consists of an olive (5Y 4/4) DIATOMACEOUS CLAY, Section 1 to 4, which grades into a pale olive (5Y 6/3) NANNOFOSSIL-RICH DIATOMACEOUS CLAY in Sections 5 and 6. Section 1 to 4 contain faint 1 cm thick pale olive (5Y 6/3) NANNOFOSSIL-RICH DIATOMACEOUS CLAY mottles.
2									SS	
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										
26										
27										
28										
29										
30										
31										
32										
33										
34										
35										
36										
37										
38										
39										
40										
41										
42										
43										
44										
45										
46										
47										
48										
49										
50										
51										
52										
53										
54										
55										
56										
57										
58										
59										
60										
61										
62										
63										
64										
65										
66										
67										
68										
69										
70										
71										
72										
73										
74										
75										
76										
77										
78										
79										
80										
81										
82										
83										
84										
85										
86										
87										
88										
89										
90										
91										
92										
93										
94										
95										
96										
97										
98										
99										
100										
101										
102										
103										
104										
105										
106										
107										
108										
109										
110										
111										
112										
113										
114										
115										
116										
117										
118										
119										
120										
121										
122										
123										
124										
125										
126										
127										
128										
129										
130										
131										
132										
133										
134										
135										
136										
137										
138										
139										
140										
141										
142										
143										
144										
145										
146										
147										
148										
149										
150										
151										
152										
153										
154										
155										
156										
157										
158										
159										
160										
161										
162										
163										
164										
165										
166										
167										
168										
169										
170										
171										
172										
173										
174										
175										



1084A-40X 354.8-364.4 mbsf										
Leg 175 Site 1084 Hole A Core 40X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	NANNOFOSSIL-RICH DIATOMACEOUS CLAY and NANNOFOSSIL-BEARING DIATOM-RICH CLAY The core consists of olive (5Y 4/3 and 5Y 5/3) NANNOFOSSIL-RICH DIATOMACEOUS CLAY in Sections 1 to 4 and olive (5Y 4/3) NANNOFOSSIL-BEARING DIATOM- RICH CLAY in Sections 5 to the end of core.
2									SS	
3										
4										
5									SS	
6										
7									PAL	




1084A-41X 364.4-374.1 mbsf										
Leg 175 Site 1084 Hole A Core 41X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								↕	SS	NANNOFOSSIL-RICH DIATOMACEOUS CLAY, DIATOMACEOUS CLAY, and NANNOFOSSIL DIATOMACEOUS CLAY The core consists of olive (5Y 4/3) NANNOFOSSIL-RICH DIATOMACEOUS CLAY in the upper part of Section 1. End of Section 1, Section 2, and Section 3, 0-50 cm, are dark olive gray (5Y 3/2) DIATOMACEOUS CLAY. Sections 3, 50 cm, to the end of the core are olive (5Y 5/4) NANNOFOSSIL DIATOMACEOUS CLAY.
2									SS	
3									IW	
4									SS	
5								↕		
6								↕	SS	
7										
8									PAL	



1084A-42X 374.1-383.7 mbsf									
Leg 175 Site 1084 Hole A Core 42X									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									SS
2									
3									
4									XRD
4									SS
4									SS
6									
5									
6									SS
8									PAL

1084A-42X


1084A-43X 383.7-393.3 mbsf										
Leg 175 Site 1084 Hole A Core 43X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	FORAMINIFER-BEARING NANNOFOSSIL-AND DIATOM-RICH CLAY, DIATOMACEOUS CLAY, and NANNOFOSSIL- AND DIATOM-RICH CLAY The core consists of olive (5Y 4/3) FORAMINIFER-BEARING NANNOFOSSIL-AND DIATOM-RICH CLAY in Sections 1 and 2. Sections 3, 4, and 5, 90 cm, are dark olive gray (5Y 3/2) DIATOMACEOUS CLAY. At Section 5, 90 cm, a sharp color change into olive (5Y 4/3) corresponds to the transition into NANNOFOSSIL- AND DIATOM-RICH CLAY to the end of the core.
2										
3									SS	
4										
5									SS	
6										
7									SS	
8										
10									PAL	



1084A-44X 393.3-402.9 mbsf										
Leg 175 Site 1084 Hole A Core 44X										
MEIRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										CLAY-RICH NANNOFOSSIL DIATOM OOZE
2									SS	The core consists of olive (5Y 5/3) CLAY-RICH NANNOFOSSIL DIATOM OOZE. The entire core has fine fractures every 0.5 cm.
3									IW	
4									SS	
5										
6										
7										
8									PAL	




1084A-45X 402.9-412.6 mbsf									
Leg 175 Site 1084 Hole A Core 45X									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									
2									
3									
4									
5									
6									
7									
8									
9									
10									




1084A-46X 412.6-422.2 mbsf										
Leg 175 Site 1084 Hole A Core 46X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										CLAY-RICH NANNOFOSSIL DIATOM OOZE, NANNOFOSSIL-BEARING DIATOM OOZE, and CLAY-BEARING NANNOFOSSIL DIATOM OOZE The core consists of an olive gray (5Y 3/2) CLAY-RICH NANNOFOSSIL DIATOM OOZE in Section 2 and Section 3, 70-150 cm, and a black (5Y 2.5/2) NANNOFOSSIL-BEARING DIATOM OOZE in Section 3, 0-70 cm. Section 4 to the end of the core is pale olive (5Y 6/3) CLAY-BEARING NANNOFOSSIL DIATOM OOZE. Core disturbance is present in Section 1.
2									SS	
3									SS	
4										
5									SS	
6										
7										
8									PAL	



1084A-46X

1084A-47X 422.2-431.9 mbsf										
Leg 175 Site 1084 Hole A Core 47X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										CLAY-BEARING DIATOMACEOUS NANNOFOSSIL OOZE and CLAY-RICH NANNOFOSSIL DIATOM OOZE The core consists of pale olive (5Y 6/3) CLAY-BEARING DIATOMACEOUS NANNOFOSSIL OOZE in Section 2 and Section 3, 0-90 cm, and an olive (5Y 4/3) CLAY-RICH NANNOFOSSIL DIATOM OOZE in Section 3, 90 cm, to the end of the core. Core disturbance is present in Section 1.
2									SS	
3										
4									SS	
5									IW	
6										
7									PAL	



1084A-48X 431.9-441.6 mbsf										
Leg 175 Site 1084 Hole A Core 48X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 4 5 6 7									SS SS PAL	NANNOFOSSIL DIATOMACEOUS CLAY and CLAY-RICH DIATOMACEOUS NANNOFOSSIL OOZE The core consists of an olive (5Y 4/3) NANNOFOSSIL DIATOMACEOUS CLAY and CLAY-RICH DIATOMACEOUS NANNOFOSSIL OOZE. The core has fine fractures (0.5 cm thick) throughout, with more severe disturbance in Section 3, 103-120 cm, and Section 5, 95-100 cm. Carbonate concretions are present in the core catcher.




1084A-49X 441.6-451.2 mbsf										
Leg 175 Site 1084 Hole A Core 49X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5 6 7										DIATOM-BEARING NANNOFOSSIL-RICH CLAY and DIATOM- AND NANNOFOSSIL-RICH CLAY The core consists of an olive (5Y 4/4, 5Y 5/3) DIATOM-BEARING NANNOFOSSIL-RICH CLAY and DIATOM- AND NANNOFOSSIL-RICH CLAY. A color change occurs over one meter in Section 4. Fine fractures occur throughout.


1084A-49X

1084A-50X 451.2-460.8 mbsf										
Leg 175 Site 1084 Hole A Core 50X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM- AND NANNOFOSSIL-BEARING CLAY, DIATOM-BEARING NANNOFOSSIL-RICH CLAY and DIATOM- AND CLAY-RICH NANNOFOSSIL OOZE The core consists of olive (5Y 5/3) DIATOM-BEARING NANNOFOSSIL- RICH CLAY in Sections 1-4, with a very dark grayish brown (2.5Y 3/2) organic carbon rich interval of DIATOM- AND NANNOFOSSIL-BEARING CLAY in Section 1, 100 cm, to Section 2, 60 cm. A pale yellow (5Y 7/3) DIATOM- AND CLAY-RICH NANNOFOSSIL OOZE extends from Section 5 to the bottom of the core. The color transition from olive to pale yellow occurs over about 80 cm. Fine fractures occur throughout the core.
2										
3										
4										
5										
6										
7										
8										
9										




1084A-51X 460.8-470.0 mbsf										
Leg 175 Site 1084 Hole A Core 51X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-BEARING CLAY-RICH NANNOFOSSIL OOZE, DIATOM-RICH NANNOFOSSIL OOZE, DIATOM-BEARING CLAY The core consists of DIATOM-BEARING CLAY-RICH NANNOFOSSIL OOZE, NANNOFOSSIL-RICH CLAY, DIATOM-BEARING CLAY. Biogenic abundances do not appear directly correlated with the color changes. Sections 1, 2, and 5 are olive (5Y 5/3), Sections 3 and 4 are very dark grayish brown, and Section 6 is pale olive (5Y 6/3). Several slightly lithified concretions occur in Section 1, 70 cm, Section 2, 70 cm, Section 4, 50 and 100 cm, and Section 6, 70 cm.
2									SS	
3										
4										
5									SS	
6									SS	
7									PAL	




1084A-52X 470.0-479.7 mbsf										
Leg 175 Site 1084 Hole A Core 52X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								///	— SS	DIATOM-RICH NANNOFOSSIL OOZE The core consists of an olive (5Y 5/3) DIATOM-RICH NANNOFOSSIL OOZE.
2							///			
3							///	— SS		
4							///			
5							///	— SS		
6							///	— PAL		


1084A-52X



1084A-53X 479.7-489.4 mbsf										
Leg 175 Site 1084 Hole A Core 53X										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL OOZE, DIATOM-BEARING NANNOFOSSIL CLAY and DIATOM-RICH NANNOFOSSIL CLAY The core consists of intervals of olive (5Y 5/3) DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL OOZE, olive gray (5Y 4/2) DIATOM-BEARING NANNOFOSSIL-RICH CLAY, and pale olive (5Y 6/3) DIATOM-RICH NANNOFOSSIL OOZE. The olive interval extends from Section 1, 0 cm, to Section 3, 110 cm. The olive gray interval extends from Section 3, 110 cm, to Section 5, 36 cm. The pale olive interval extends from Section 5, 36 cm to the end of the core. The transition from the olive to olive gray interval is gradational over 40 cm. Sections 1, 5, and 6 were dropped.
2									SS	
3									IW	
4									SS	
5									SS	
6									PAL	


1084A-53X

1084A-54X 489.4-499.1 mbsf										
Leg 175 Site 1084 Hole A Core 54X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5 6										DIATOM- AND NANNOFOSSIL-BEARING CLAY, DIATOM-BEARING NANNOFOSSIL CLAY, and DIATOM-BEARING NANNOFOSSIL-RICH CLAY The core consists of light yellowish brown (5Y 6/3) DIATOM-BEARING NANNOFOSSIL CLAY and DIATOM-BEARING NANNOFOSSIL-RICH CLAY with an interval of olive brown (5Y 4/3) DIATOM- AND NANNOFOSSIL-BEARING CLAY in Section 2, 70 cm, to Section 3, 30 cm. The transition between colors occurs over 30 cm on the top and the bottom of the dark interval. Section 1 was severely disturbed. SS SS PAL









1084A-58X 528.0-537.7 mbsf										
Leg 175 Site 1084 Hole A Core 58X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING NANNOFOSSIL CLAY and NANNOFOSSIL CLAY The core consists of olive gray (5Y 5/2) NANNOFOSSIL CLAY and olive (5Y 5/3 and 5Y 4/3) DIATOM-BEARING NANNOFOSSIL CLAY. Olive gray colored interval is seen from Section 3, 110 cm, to Section 4, 90 cm.
2									SS	
3										
4									SS	
5										
6									PAL	





1084A-59X 537.7-547.3 mbsf										
Leg 175 Site 1084 Hole A Core 59X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5 6										NANNOFOSSIL CLAY The core consists of olive (5Y 4/3) NANNOFOSSIL CLAY. The interval in Section 4, 20-80 cm is dark olive gray (5Y 3/2) colored. — SS — IW — SS — PAL

I084A-59X



1084A-60X 547.3-557.0 mbsf										
Leg 175 Site 1084 Hole A Core 60X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										CLAY-RICH NANNOFOSSIL OOZE, DIATOM-BEARING CLAY-RICH NANNOFOSSIL OOZE and NANNOFOSSIL OOZE. The core consists of intervals of olive gray (5Y 5/2) CLAY-RICH NANNOFOSSIL, dark olive gray (5Y 3/2) DIATOM-BEARING CLAY-RICH NANNOFOSSIL OOZE, and pale olive (5Y 6/3) NANNOFOSSIL OOZE. The olive gray interval extends from Section 1, 0 cm, to Section 4, 109 cm. The dark olive gray interval extends from Section 4, 109 cm, to Section 5, 5 cm. The pale olive interval extends from Section 5, 5 cm, to the end of the core. The contact between the olive gray and dark olive gray interval appears gradational over 10 cm. Rip ups, 1 cm by 1 cm in diameter, may be present in the upper portion of the dark olive gray interval.
2										
3										
4										
5										
6										
7										
8										








1084A-62X 566.6-576.2 mbsf										
Leg 175 Site 1084 Hole A Core 62X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										CLAYEY NANNOFOSSIL OOZE and DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE The core consists of olive gray (5Y 5/2 and 5Y 4/2) CLAYEY NANNOFOSSIL OOZE. Dark olive gray (5Y 3/2) DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE intervals occur in Section 3, 110-150 cm, Section 7, 120-150 cm, and Section 8. SS SS IW SS PAL
2										
3										
4										
5										
6										
7										
8										
9										
10										

1084A-62X



1084A-64X 585.9-595.5 mbsf										
Leg 175 Site 1084 Hole A Core 64X										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5 6 7									SS SS SS PAL	CLAY-RICH NANNOFOSSIL OOZE and CLAYEY NANNOFOSSIL OOZE The core consists of olive gray (5Y 4/2) CLAYEY NANNOFOSSIL OOZE. A light olive gray (5Y 6/2) CLAY-RICH NANNOFOSSIL OOZE interval occurs in Section 2, 0-50 cm.


1084A-64X




1084B-1H 0.0-4.1 mbsf										
Leg 175 Site 1084 Hole B Core 1H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								ooo	SS	DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL CLAY and DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL-RICH CLAY The core consists of olive (5Y 5/3) DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL CLAY and olive (5Y 4/3) DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL-RICH CLAY.
2									SS	
3										
4										




1084B-2H 4.1-13.6 mbsf										
Leg 175 Site 1084 Hole B Core 2H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	FORAMINIFER-BEARING DIATOM- AND NANNOFOSSIL-RICH CLAY The core consists of dark olive gray (5Y 3/2), olive gray (5Y 4/2), and olive (5Y 4/3) FORAMINIFER-BEARING DIATOM- AND NANNOFOSSIL-RICH CLAY. Sections 1, 2, and 3 are dark olive gray, Section 4 is olive gray, and Sections 5 and 6 are olive. The color changes are gradual.
2										
3									SS	
4										
5									SS	
6										



1084B-3H 13.6-23.1 mbsf										
Leg 175 Site 1084 Hole B Core 3H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	FORAMINIFER- AND NANNOFOSSIL-BEARING CLAY and FORAMINIFER-BEARING NANNOFOSSIL- AND DIATOM-RICH CLAY and FORAMINIFER- AND NANNOFOSSIL-BEARING DIATOM-RICH CLAY, and DIATOM-BEARING FORAMINIFER-RICH NANNOFOSSIL CLAY The core consists of dark olive gray (5Y 3/2) FORAMINIFER- AND NANNOFOSSIL-BEARING CLAY, olive gray (5Y 4/2) FORAMINIFER-BEARING NANNOFOSSIL- AND DIATOM-RICH CLAY, olive (5Y 4/3) DIATOM-BEARING FORAMINIFER-RICH NANNOFOSSIL CLAY, and black (5Y 2.5/2) FORAMINIFER- AND NANNOFOSSIL-BEARING DIATOM-RICH CLAY.
2										
3									SS	
4									SS	
5									SS	
6									SS	
7										

1084B-3H


1084B-4H 23.1-32.6 mbsf										
Leg 175 Site 1084 Hole B Core 4H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								SS		DIATOM- and NANNOFOSSIL-RICH CLAY and CLAYEY NANNOFOSSIL OOZE The core consists of dark olive gray (5Y 3/2) DIATOM- and NANNOFOSSIL-RICH CLAY with abundant diatom resting spores and olive (5Y 4/3) and olive gray (5Y 4/2) CLAYEY NANNOFOSSIL OOZE.
2								SS		
3								SS		
4								SS		
5								SS		
6										
7										
8										
9										
10										



1084B-5H 32.6-42.1 mbsf										
Leg 175 Site 1084 Hole B Core 5H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	NANNOFOSSIL-BEARING DIATOM-RICH CLAY, DIATOM-RICH NANNOFOSSIL CLAY and DIATOM-BEARING CLAY The core consists of black (5Y 2.5/1) NANNOFOSSIL-BEARING DIATOM-RICH CLAY and olive (5Y 4/3) DIATOM-RICH NANNOFOSSIL CLAY and black (2.5Y 2.5/1) DIATOM-BEARING CLAY. Only the upper boundaries of the DIATOM-RICH NANNOFOSSIL CLAY show large burrow traces.
2										
3										
4										
5										
6										
7										
8										
9										
10										


1084B-5H


1084B-6H 42.1-51.6 mbsf								
Leg 175 Site 1084 Hole B Core 6H								
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	
								DESCRIPTION
1								DIATOM-RICH CLAY, NANNOFOSSIL- AND DIATOM-RICH CLAY, DIATOM- AND FORAMINIFER-RICH NANNOFOSSIL CLAY, and DIATOM- AND FORAMINIFER-RICH NANNOFOSSIL OOZE The core consists of black (5Y 2.5/2) DIATOM-RICH CLAY, olive (5Y 4/3) NANNOFOSIL- AND DIATOM-RICH CLAY, olive (5Y 4/2) FORAMINIFER-RICH NANNOFOSSIL CLAY, and olive (5Y 4/3) DIATOM- AND FORAMINIFER-RICH NANNOFOSSIL OOZE. Changes in lithology are gradual.
2								
3								
4								
5								
6								SS
7								
8								
9								
10								



1084B-7H 51.6-61.1 mbsf										
Leg 175 Site 1084 Hole B Core 7H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	FORAMINIFER-BEARING CLAYEY NANNOFOSSIL OOZE, DIATOM-BEARING NANNOFOSSIL-RICH CLAY, and DIATOM-RICH CLAYEY NANNOFOSSIL OOZE The core consists of dark olive gray (5Y 3/2) FORAMINIFER-BEARING CLAYEY NANNOFOSSIL OOZE, black (5Y 2.5/2) DIATOM-BEARING NANNOFOSSIL-RICH CLAY, and light olive gray (5Y 6/2) and olive gray (5Y 5/2) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE. Burrow traces occur at the upper boundary of the black DIATOM-BEARING NANNOFOSSIL-RICH CLAY and the DIATOM-RICH CLAYEY NANNOFOSSIL OOZE.
2										
3									SS	
4										
5										
6									SS	
7										
8										
9										
10										


1084B-7H


1084B-8H 61.1-70.6 mbsf										
Leg 175 Site 1084 Hole B Core 8H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										FORAMINIFER-BEARING CLAYEY NANNOFOSSIL OOZE and DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE The core consists of olive gray (5Y 5/2) FORAMINIFER-BEARING CLAYEY NANNOFOSSIL OOZE and olive (5Y 5/3) DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE. The olive gray interval extends from Section 1, 0 cm, to Section 2, 20 cm. Burrows range in diameter from 1 to 2 cm.
2									SS	
3										
4										
5									SS	
6										



1084B-9H 70.6-80.1 mbsf										
Leg 175 Site 1084 Hole B Core 9H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										FORAMINIFER-BEARING CLAYEY NANNOFOSSIL OOZE, DIATOM-BEARING NANNOFOSSIL OOZE, FORAMINIFER-BEARING CLAY-RICH NANNOFOSSIL OOZE The core consists of intercalated intervals of olive (5Y 4/3) FORAMINIFER-BEARING CLAYEY NANNOFOSSIL OOZE (Section 1, 0 cm, to Section 2, 10 cm), light olive gray (5Y 6/2) DIATOM-BEARING NANNOFOSSIL OOZE (Section 2, 10 cm, to Section 3, 10 cm), and olive (5Y 5/3) FORAMINIFER-BEARING CLAY-RICH NANNOFOSSIL OOZE (Section 3, 10 cm to the end of the core).
2										
3										
4										
5										
6										
7										
8										
9										
10										



1084B-10H 80.1-89.6 mbsf									
Leg 175 Site 1084 Hole B Core 10H									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									
2									SS
3									
4									
5									
6									SS
7									
8									
\ CLAYEY NANNOFOSSIL OOZE and NANNOFOSSIL-RICH CLAY									
The core consists of olive (5Y 4/3) CLAYEY NANNOFOSSIL OOZE with an interval of dark olive gray (5Y 3/2) NANNOFOSSIL-RICH CLAY. This interval extends from Section 1, 90 cm, to Section 2, 90 cm. Its contacts with the olive CLAYEY NANNOFOSSIL OOZE are gradational over 40 cm.									




1084B-11H 89.6-99.1 mbsf										
Leg 175 Site 1084 Hole B Core 11H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	NANNOFOSSIL-RICH CLAY and NANNOFOSSIL-RICH CLAY The core consists of olive (5Y 5/3) NANNOFOSSIL-RICH CLAY with intercalated intervals of black (5Y 2.5/2) NANNOFOSSIL-RICH CLAY. Black intervals extend from Section 2, 80 cm, to Section 3, 20 cm, and Section 3, 120 cm, to the end of the core. Intervals have gradational contacts which grade over 30-40 cm.
2								///		
3								↑ /// ↓		
4										
5										
6								~	SS	




1084B-12H 99.1-108.6 mbsf									
Leg 175 Site 1084 Hole B Core 12H									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1	1								
2	2								
3	3								
4	4								
5	5								
6	6								
7	7								
8	8								
10									

FORAMINIFER-BEARING NANNOFOSSIL-RICH CLAY, CLAY, and NANNOFOSSIL-BEARING CLAY


The core contains an olive (5Y 4/3) CLAY with shorter intervals of a black (5Y 2.5/2) FORAMINIFER-BEARING NANNOFOSSIL-RICH CLAY in Section 1 and a dark olive gray (5Y 3/2) NANNOFOSSIL-BEARING CLAY in Section 5. Severe core disturbance is present in the lower portion of Section 2.





1084B-14H 118.1-127.6 mbsf									
Leg 175 Site 1084 Hole B Core 14H									
METERS	SECTION	GRAPHIC LITH.	BIO TURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									
2									
3									
4									
5									
6									
7									
8									
9									
10									

I084B-14H





1084B-16H 137.1-146.6 mbsf										
Leg 175 Site 1084 Hole B Core 16H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING NANNOFOSSIL CLAY, DIATOM- AND NANNOFOSSIL-BEARING CLAY and NANNOFOSSIL OOZE The core consists of olive (5Y 4/3) DIATOM-BEARING NANNOFOSSIL CLAY (Section 1, 0 cm, to Section 6, 97 cm, and Section 6, 140 cm, to Section 7, 80 cm), black (5Y 2.5/2) DIATOM- AND NANNOFOSSIL-BEARING CLAY (Section 6, 140 cm, to Section 7, 80 cm), and pale olive (5Y 6/3) NANNOFOSSIL OOZE (Section 7, 80 cm to the end of the core).
2										
3									SS	
4										
5										
6									SS	
7									SS	
8										
9									PAL	

I084B-16H


1084B-17H 146.6-156.1 mbsf										
Leg 175 Site 1084 Hole B Core 17H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-RICH NANNOFOSSIL CLAY The core consists of olive (5Y 4/3 and 5Y 5/3) DIATOM-RICH NANNOFOSSIL CLAY. Gas expansion voids are common in the upper part of the core.
2										
3									SS	
4										
5										
6									SS	
7										
8									PAL	
9										


1084B-17H



1084B-19H 165.6-175.1 mbsf										
Leg 175 Site 1084 Hole B Core 19H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL CLAY The core consists of olive (5Y 4/3) DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL CLAY.
2										
3										
4									SS	
5										
6									SS	
7										
8									PAL	


1084B-19H


1084B-20H 175.1-182.8 mbsf										
Leg 175 Site 1084 Hole B Core 20H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 5 6 7									SS SS PAL	DIATOM-BEARING NANNOFOSSIL CLAY and NANNOFOSSIL CLAY The core consists of olive (5Y 4/3) DIATOM-BEARING NANNOFOSSIL CLAY and black (5Y 2.5/1) NANNOFOSSIL CLAY. The transitions between different colored intervals occur in Section 2, 70-100 cm, and Section 6, 50 cm, to the top 20 cm of the core catcher. Bioturbation is evident in olive colored sections.








1084C-1H 0.0-8.1 mbsf						
Leg 175 Site 1084 Hole C Core 1H						
METERS	SECTION GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO. FOSSILS	DISTURB. SAMPLE
						DESCRIPTION
0						~ DIATOM-BEARING CLAYEY NANNOFOSSIL OOOZE and DIATOM- AND NANNOFOSSIL-BEARING CLAY The core consists of olive (5Y 5/3) DIATOM-BEARING CLAYEY NANNOFOSSIL OOOZE (Section 1) and a dark olive gray (5Y 3/2) DIATOM- AND NANNOFOSSIL-BEARING CLAY.
1					— SS	
2						
3						
4						
5					— SS	
6						
7						
8						


1084C-1H


1084C-2H 8.1-17.6 mbsf										
Leg 175 Site 1084 Hole C Core 2H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										NANNOFOSSIL-RICH DIATOMACEOUS CLAY and DIATOM-RICH CLAYEY NANNOFOSSIL OOZE The core consists of dark olive gray (5Y 3/2) NANNOFOSSIL-RICH DIATOMACEOUS CLAY. Section 6, 40 cm, to the end of the core is olive (5Y 4/3) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE.
2								SS		
3										
4										
5										
6										
7									SS	



1084C-3H 17.6-27.1 mbsf										
Leg 175 Site 1084 Hole C Core 3H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM- AND NANNOFOSSIL-BEARING CLAY, DIATOM BEARING NANNOFOSSIL CLAY, DIATOM-RICH NANNOFOSSIL CLAY, and DIATOM-RICH CLAYEY NANNOFOSSIL OOZE. The core consists of very dark gray (5Y 3/1) DIATOM- AND NANNOFOSSIL-BEARING CLAY (Section 1 to Section 2, 50 cm) olive gray DIATOM BEARING NANNOFOSSIL CLAY (Section 2, 50 cm, to Section 3, 80 cm), very dark gray (5Y 3/1) DIATOM- AND NANNOFOSSIL-BEARING CLAY (Section 3, 80 cm, to Section 4, 50 cm), olive (5Y 5/4) DIATOM-RICH NANNOFOSSIL CLAY (Section 4, 50 cm, to Section 5, 50 cm) olive gray (5Y 4/2) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE (Section 5, 50 cm, to the end of the core).
2									SS	
3										
4									SS	
5									SS	




1084C-4H 27.1-36.6 mbsf										
Leg 175 Site 1084 Hole C Core 4H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1 2 3 4 6 5 8 6 7 10									—SS —SS	~ NANNOFOSSIL DIATOMACEOUS CLAY and DIATOMACEOUS CLAY The core consists of olive gray (5Y 4/2) NANNOFOSSIL DIATOMACEOUS CLAY and black (5Y 2.5/1) DIATOMACEOUS CLAY.



1084C-5H 36.6-46.1 mbsf										
Leg 175 Site 1084 Hole C Core 5H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	NANNOFOSSIL-BEARING CLAY, DIATOM-RICH CLAYEY NANNOFOSSIL OOZE, and NANNOFOSSIL-BEARING DIATOMACEOUS CLAY The core consists of black (5Y 2.5/1) NANNOFOSSIL-BEARING CLAY (Section 1, 0-100 cm), pale olive (5Y 6/3) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE (Section 1, 100 cm, to Section 2, 75 cm), dark olive gray (5Y 3/2) NANNOFOSSIL-BEARING DIATOMACEOUS CLAY (Section 2, 75 cm, to Section 3, 100 cm), black (5Y 2.5/1) NANNOFOSSIL-BEARING CLAY (Section 3, 100 cm to Section 4, 60 cm), pale olive (5Y 6/3) DIATOM-RICH CLAYEY NANNOFOSSIL OOZE (Section 4, 60 cm, to Section 5, 30 cm), and dark olive gray (5Y 3/2) NANNOFOSSIL-BEARING DIATOMACEOUS CLAY (Section 5, 30 cm, to the end of the core).
2									SS	
3										
4										
5										
6									SS	
7										
8										
9										
10										




1084C-6H 46.1-55.6 mbsf										
Leg 175 Site 1084 Hole C Core 6H										
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE and DIATOM-BEARING NANNOFOSSIL-RICH CLAY The core consists of DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE and DIATOM-BEARING NANNOFOSSIL-RICH CLAY. Sections 1 to 4 are olive (5Y 5/3) DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE. Section 4 contains black (5Y 2.5/1) 3 cm thick mottles. Section 5, 0-100 cm, is light olive gray (5Y 6/2) DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE. Section 5, 100 cm, to Section 6, 150 cm, is olive (5Y 5/3) DIATOM-BEARING CLAYEY NANNOFOSSIL OOZE. Section 7 to the end of the core is black (5Y 2.5/1) DIATOM-BEARING NANNOFOSSIL-RICH CLAY.
2									SS	
3										
4										
5									SS	
6										
7									SS	



1084C-7H 55.6-65.1 mbsf								
Leg 175 Site 1084 Hole C Core 7H								
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DESCRIPTION
1								NANNOFOSSIL CLAY and DIATOM-RICH NANNOFOSSIL OOZE and DIATOM-RICH NANNOFOSSIL CLAY The core consists of black (5Y 3/1) NANNOFOSSIL CLAY, pale olive (5Y 6/3) DIATOM-RICH NANNOFOSSIL OOZE, and olive (5Y 4/3) DIATOM-RICH NANNOFOSSIL CLAY.
2								
3								
4								
5								
6								NANNOFOSSIL CLAY and DIATOM-RICH NANNOFOSSIL OOZE and DIATOM-RICH NANNOFOSSIL CLAY The core consists of black (5Y 3/1) NANNOFOSSIL CLAY, pale olive (5Y 6/3) DIATOM-RICH NANNOFOSSIL OOZE, and olive (5Y 4/3) DIATOM-RICH NANNOFOSSIL CLAY.
7								
8								
9								
10								




1084C-8H 65.1-74.6 mbsf								
Leg 175 Site 1084 Hole C Core 8H								
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	
								DESCRIPTION
1								DIATOM-BEARING NANNOFOSSIL OOZE, DIATOM-BEARING NANNOFOSSIL CLAY, DIATOM-BEARING NANNOFOSSIL CLAY, and DIATOM-RICH NANNOFOSSIL OOZE The core consists of olive (5Y 4/3) DIATOM-BEARING NANNOFOSSIL CLAY, olive (5Y 5/3) DIATOM-BEARING NANNOFOSSIL OOZE, and pale olive (5Y 6/3) DIATOM-RICH NANNOFOSSIL OOZE.
2								
3								
4								
5								
6								
7								
8								
9								
10								




1084C-9H 74.6-84.1 mbsf											
Leg 175 Site 1084 Hole C Core 9H											
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION	
1										DIATOM-RICH NANNOFOSSIL OOZE, DIATOM-BEARING NANNOFOSSIL OOZE, and NANNOFOSSIL CLAY The core consists of olive gray (5Y 5/2) DIATOM-RICH NANNOFOSSIL OOZE, olive (5Y 4/3) DIATOM-BEARING NANNOFOSSIL OOZE, and very dark gray (5Y 3/2) NANNOFOSSIL CLAY.	
2											
3											
4											
5											
6									SS		
7									SS		
8											
9											
10									SS		



1084C-10H 84.1-93.6 mbsf								
Leg 175 Site 1084 Hole C Core 10H								
METRES	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	
								DESCRIPTION
1								DIATOM-RICH NANNOFOSSIL OOZE The core consists of dark olive gray (5Y 3/2) to olive gray (5Y 4/2) DIATOM-RICH NANNOFOSSIL OOZE. Brown oxidation rims and veins occur on the outside of the sediment in Sections 5, 6, and 7.
2								
3								
4								
5								
6								
7								
8								
9								
10								




1084C-11H 93.6-103.1 mbsf										
Leg 175 Site 1084 Hole C Core 11H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1									SS	DIATOM-RICH NANNOFOSSIL CLAY, DIATOM- AND NANNOFOSSIL-RICH CLAY, DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL-RICH CLAY The core consists of black (5Y 2.5/1) DIATOM-RICH NANNOFOSSIL CLAY, DIATOM- AND NANNOFOSSIL-RICH CLAY, and black (2.5Y 2.5/1) DIATOM- AND FORAMINIFER-BEARING NANNOFOSSIL-RICH CLAY.
2									SS	
3										
4										
5										
6										
7									SS	





1084C-12H 103.1-112.6 mbsf									
Leg 175 Site 1084 Hole C Core 12H									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									SS
2									
3									
4									
5									SS
6									
7									
8									
9									
10									SS


DIATOM-RICH CLAY and DIATOM- AND NANNOFOSSIL-RICH CLAY

The core consists of black (2.5Y 2.5/1) DIATOM-RICH CLAY and olive (5Y 4/3) DIATOM- AND NANNOFOSSIL-RICH CLAY. A gradual transition from DIATOM-RICH CLAY to DIATOM- AND NANNOFOSSIL-RICH CLAY is present in Sections 3 and 4.






1084C-14H 122.1-131.6 mbsf										
Leg 175 Site 1084 Hole C Core 14H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								↕		NANNOFOSSIL-RICH CLAY and CLAY The core consists of a dark olive (5Y 3/2) and olive (5Y 4/3) NANNOFOSSIL-RICH CLAY with an interval of very dark gray (5Y 3/1) CLAY. Gas expansion voids are present throughout the core.
2								↕↕↕↕		
3										
4										
5										
6										
7								↕		
8								↕		
9										
10										




1084C-15H 131.6-141.1 mbsf										
Leg 175 Site 1084 Hole C Core 15H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								~ ~		NANNOFOSSIL-RICH CLAY and NANNOFOSSIL-BEARING CLAY The core consists of olive gray (5Y 4/2) NANNOFOSSIL-RICH CLAY which grades to a dark olive gray (5Y 3/2) NANNOFOSSIL-BEARING CLAY by Section 5.
2								↑ ↓		
3								↑ ↓		
4								↑ ↓		
5								↑ ↓		
6								↑ ↓		
7								↑ ↓		
8								↑ ↓		
9								↑ ↓		
10								↑ ↓		




1084C-16H 141.1-150.6 mbsf										
Leg 175 Site 1084 Hole C Core 16H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										NANNOFOSSIL-BEARING CLAY and DIATOM-BEARING NANNOFOSSIL CLAY The core consists of black (5Y 2.5/2) NANNOFOSSIL-BEARING CLAY and olive DIATOM-BEARING NANNOFOSSIL CLAY. Color changes occur gradually over 20 to 60 cm. Bioturbation is most evident at color transitions within the core.
2										
3										
4										
5										
6										
7										
8										
9										
10										






1084C-17H 150.6-160.1 mbsf										
Leg 175 Site 1084 Hole C Core 17H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHNO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1										DIATOM-BEARING NANNOFOSSIL CLAY The core consists of olive (5Y4/3 and 5Y 5/3) DIATOM-BEARING NANNOFOSSIL CLAY. Dark olive (5Y 3/2) colored sediments are only present in Section 5, 0-40 cm, and 85 cm to Section 6, 20 cm. Gas expansion voids are common throughout the core.
2										
3										
4										
5										
6										
7										
8										
9										
10										



1084C-18H 160.1-169.6 mbsf										
Leg 175 Site 1084 Hole C Core 18H										
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE	DESCRIPTION
1								↕		DIATOM-BEARING NANNOFOSSIL CLAY and FORAMINIFER- AND DIATOM-BEARING NANNOFOSSIL CLAY The core consists of dark olive gray (5Y 3/2) DIATOM-BEARING NANNOFOSSIL CLAY, and olive (5Y 5/3 and 5Y 4/3) colored FORAMINIFER- AND DIATOM-BEARING NANNOFOSSIL CLAY (Section 4, 10-80 cm, and Section 5, 125 cm, to the end of the core).
2								↕		
3								↕		
4								— SS		
5								↕		
6								↕		
7								— SS		







1084C-21H 188.6-198.1 mbsf									
Leg 175 Site 1084 Hole C Core 21H									
METERS	SECTION	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	ICHO.	FOSSILS	DISTURB.	SAMPLE
DESCRIPTION									
1									
2									SS
3									
4									
5									
6									
7									SS
~ NANNOFOSSIL CLAY and DIATOM-RICH NANNOFOSSIL CLAY The core consists of olive NANNOFOSSIL CLAY and dark olive gray (5Y 3/2) DIATOM-RICH NANNOFOSSIL CLAY.									



